

**State of California
The Resources Agency
DEPARTMENT OF FISH AND GAME**

**2000-2001 ANNUAL REPORT
TRINITY RIVER TRIBUTARIES
WINTER-RUN STEELHEAD SPAWNING SURVEY REPORT
PROJECT 1d1**

Prepared by

**Patrick Garrison
Northern California - North Coast Region**

**Steelhead Research and Monitoring Program
January 2002**

ABSTRACT

Spawning surveys for winter-run steelhead were conducted on tributaries of the Trinity River from March 1 through May 31, 2001. We walked a total of 14 tributaries, three or more times each, for a single pass total of 95.5 km. Only 29 adult steelhead were observed, and 37 redds were marked and recorded. The highest density of steelhead redds occurred in Eltapom Creek at 2.4 redds per km, while the lowest density occurred in four different tributaries where no redds were observed. Tributaries to be surveyed were randomly selected from a universe of all anadromous Trinity River tributaries upstream of the New River. Coincidentally, several selected tributaries duplicate the work of D.A. LaFaunce (1964), D.W. Rogers (1971, 1972), Ed Miller (1974), and the South Fork Steelhead Monitoring Project (1990-1995), and all applicable comparisons are made.

Introduction

The current state of knowledge regarding steelhead spawning habitat is limited. Most prior spawner surveys within the KMP ESU concentrated on salmon and were therefore terminated prior to steelhead spawning. Prior surveys have been conducted on main-stem Trinity tributaries in 1964, 1971, 1972, and 1974 to monitor the effect of Lewiston Dam on steelhead populations. Most recently, steelhead spawning surveys were conducted in South Fork Trinity River tributaries in 1989 - 1995 under the Department's Trinity River Project. Basin-wide estimates of steelhead abundance provide little information on steelhead spawning. Information regarding spawning habitat will help to assess this critical component of steelhead life history.


Study Area

The area covered by these spawning surveys includes all anadromous tributaries of the Trinity basin upstream of the New River, including the South Fork of the Trinity River. A stratified random sampling design was used to select tributaries within the basin. To develop a sampling universe, all anadromous tributaries within the named basins had to be identified. The entire basin was then stratified into two sub-basins, the South Fork and the main-stem, each of which was sampled approximately evenly. Originally, eight tributaries were selected from each basin. Three tributaries had to be dropped from the main-stem basin due to high flow problems. No replacement tributaries were chosen in the main-stem due to time restraints. The following Trinity River tributaries were surveyed from their confluence to a upstream migrational barrier except where noted.

Smoky Creek was surveyed from the South Fork Trinity River confluence to a waterfall barrier 6.7 km upstream. Access is only available through private property owned by Jon Ostrat near Silver Creek.

Rattlesnake Creek was surveyed from the South Fork Trinity River confluence to a waterfall barrier 16.21 km upstream. Access is available via State Route 36.

Figure 1. Map of Trinity basin with spawner survey tributaries



Plummer Creek was surveyed from the South Fork Trinity River confluence to a waterfall barrier 5.18 km upstream. Access is available through River Spirit Land Conservancy or by Friend Lake trail. CDFG currently has a land-owner access agreement with David Rose of the River Spirit Land Conservancy in which a small fee is paid to maintain their extensive private road system in exchange for year-round access.

Eltapom Creek was surveyed from the South Fork Trinity River confluence to a waterfall barrier 1.26 km upstream. Access is only available by crossing the South Fork Trinity River (SFTR), off of FH 311. A raft is recommended and sometimes necessary for crossing the SFTR at higher flows, especially in March and early April.

Pelletreau Creek was surveyed from the South Fork Trinity River confluence to a log jam/ depositional barrier 1.41km upstream. Access is made from FH 311 just south of Hyampom. Downstream from the 311 bridge (about 200 meters) was surveyed with binoculars after private property access permission was denied.

East Fork of Hayfork Creek was surveyed from its confluence with Hayfork Creek to Byron Gulch approximately 6.77 km upstream. There is no apparent barrier on E.F. Hayfork Creek; the survey end point was selected where suitable steelhead spawning habitat ended and low flows made fish access problematic. Access is available via FH 343. Extra safety precautions should be observed in the East Fork Hayfork Creek basin as three residents of the area are currently missing under suspicious circumstances.

Potato Creek was surveyed from its confluence East Fork Hayfork Creek to waterfall barrier 4.03 km upstream. Access is available via FH 343.

Tule Creek was surveyed from its confluence with Hayfork Creek to a culvert barrier created by the 31N31 road, approximately 8.41 miles upstream. The majority of Tule Creek is accessible via FH 10.

East Fork of the North Fork Trinity River (EFNFTR) was excluded from this year's survey after it was determined that size and high flows impeded proper surveying. EFNFTTR was previously surveyed from its confluence with North Fork Trinity River to an unnamed tributary above East Branch. The unnamed tributary above East Branch was selected as an end point because it was the farthest point accessible from the road within one day's hike.

Deadwood Creek was surveyed from its confluence with the Trinity River to a waterfall barrier 3.82 km upstream. Access is available from Deadwood Road. Deadwood is the uppermost tributary to the Trinity River below Lewiston Dam.

Dutch Creek was surveyed from its confluence with the Trinity River to a culvert barrier created by a SPI logging road. Access is available via Dutch Creek Road, but it is not

advisable to drive to the confluence, especially during winter flows. Access is available to the uppermost reach via an SPI logging road.

South Fork of Indian Creek was surveyed from its confluence with Indian Creek to a waterfall barrier 1.49 km upstream. Access is available Redding Creek Rd. and by an unnamed SPI logging road. Avoid the paranoid miner if possible.

Redding Creek was surveyed from its confluence with the Trinity River to Byron Gulch approximately 20.86 km upstream. No apparent barrier was encountered during the survey, which ended at a point two days hike in from any driveable access. Access is available via Redding Creek road and several SPI logging roads, the upper two sections are only accessible by hiking cross-county.

Results

Table 1. Number of live steelhead observed, steelhead redds, distance surveyed, and redds/km for Trinity River tributaries.

Creek	Distance Surveyed (km)	Redds	Redds/km	Adult Steelhead observed	Adult Steelhead/km
Tule	8.4	4	0.48	4	0.48
E.F. Hayfork	6.8	10	1.47	6	0.88
Deadwood	3.8	2	0.53	6	1.58
Eltapom	1.3	5	3.84	2	1.53
Pelletreau	1.4	0	0	0	0
S.F. Indian	1.5	0	0	0	0
Redding	20.9	8	0.38	6	0.29
Plummer*	5.2	0	0	0	0
Potato	4.0	2	0.50	0	0
Rattlesnake	16.2	5	0.31	3	0.18
Dutch	5.6	0	0	0	0
Smoky*	4.1	3	0.73	0	0
EFNF Trinity*	16.3	3	0.18	5	0.31
Totals	95.5	37	0.47	31	0.30

*Poor conditions and less numerous surveys portray fewer redds or fish than most likely were present.

Table 2. Trinity River Steelhead Spawning Survey Timing; Frequency surveyed and dates.

Creek	# of times surveyed	Dates Surveyed
Smoky	1	4/15
Rattlesnake	3	3/26-3/27, 4/24-4/25, 5/15-5/17
Eltapom	3	3/20, 4/9, 5/7
Pelletreau	3	3/20, 4/9, 5/7
E.F. Hayfork	3	2/8, 2/27, 3/20-3/21, 4/16, 5/10
Potato	3	3/8, 4/2, 4/23
Tule	4	3/1, 3/7, 4/4-4/5, 5/1, 6/5
Big French	2	4/11-4/12, 5/23-5/24
E.F.N.F. Trinity	2	2/5-2/7, 3/28
Deadwood	4	3/13, 4/2, 4/12, 5/2
Dutch	3	3/12, 4/19, 5/8, 5/29
S.F. Indian	2	3/12, 4/19
Reading	4	3/15-3/19, 4/10-4/12, 5/2-5/3, 5/22-5/23
Plummer	1	3/21

Information about each individual redd, including dimensions, habitat type, substrate, depth of water, and presence of fish was taken for all 37 redd sightings. This information is stored in Department files for future comparison. A map of each individual redd location, recorded by global positioning systems are also available in Department files.

Discussion

Redd or spawning surveys serve as a good, but partially incomplete means of monitoring steelhead spawning escapement. For future clarification, I will use the terminology “redd survey” and “spawning survey” interchangeably for the remainder of this report. Redd surveys are most appropriate when other means of estimating adult escapement or spawning success are not appropriate or impossible to conduct. In the Trinity basin, problems do occur which limit the ability to estimate of winter-run steelhead. High flows and the extended length of adult steelhead migration make weir estimates partial at best. Several weirs were constructed to estimate winter-run steelhead escapement by the Department in 1989-1992 at Sandy Bar and at Forest Glen on the South Fork of the

Trinity River (CDFG, 1990-95). Efforts were finally terminated after multiple blow-outs due to high flows.

Steelhead run size is highly variable from year to year, but the data gathered over the last 40 years in the Trinity basin shows the general trend that steelhead run sizes are diminishing. This year proves no different, with steelhead returns that are highly variable between creeks, but lower overall. Work by D.A. La Faunce in 1964 sets up good baseline numbers for natural production of steelhead in the Trinity basin. Those surveys show that steelhead adult estimates were markedly higher in 1964 than in any of the following years.

Table 3. Results of work by D.A. LaFaunce, 1964. A Steelhead Spawning Survey of the Upper Trinity River System.

Tributary surveyed	Distance surveyed (km.)	Redds observed	Redds/km.
Deadwood Creek	1.66	27	16.26
S.F. Indian Creek	0.37	4	10.8
E.F.N.F. Trinity River	12.0	218	18.16
Redding Creek	16.3	279	17.11
Dutch Creek	2.6	72	27.6

Surveys conducted by D.W. Rogers in 1971, show that there had already been a sharp decline in steelhead spawning since 1964. This could have been an anomalous year, but surveys in 1972-1974 of other tributaries in the system continue to show this general trend.

Table 4. Results of steelhead spawning surveys conducted by D.W. Rogers (1971).

Tributary surveyed	Distance surveyed (km.)	Redds observed	Redds/km.
Deadwood Creek	3.7	0	0
S.F. Indian Creek	1.85	3	1.62
Redding Creek	19.25	35	1.81
Dutch Creek	1.85	0	0

Table 5. Results of steelhead spawning surveys conducted by Ed Miller (1974)

Tributary surveyed	Distance surveyed (km.)	Redds observed	Redds/km.
Tule Creek	4.25	12	2.82

The South Fork Steelhead Monitoring Project conducted steelhead spawning surveys in the South Fork Trinity basin from 1990-1995 in conjunction with adult steelhead electrofishing to monitor spawning success (unsuccessful). These surveys are the most complete data-set to date, showing major spawning trends of steelhead in the Trinity basin. Eltapom creek is the jewel of the South Fork, consistently having the highest redd densities in the basin. There is great variation in redd densities from year to year; high quality tributaries fluctuate from two to 15 redd/km each year, while poor quality tributaries fluctuate between zero and two redds/km each year. I believe this year's results again most closely resemble returns from 1992-1993. These were low return years, which were immediately followed by a year of higher adult returns.

Table 6. Results of work by CDFG (1990-1995) Steelhead Spawning Survey, compared with Current Spawning Survey (2000). Distance surveyed and redd densities for 1990-1995, and 2000.

Tributary surveyed	Distance surveyed (km.) ¹	Redds s/ km 1990	Redds/ km 1991	Redds/ km 1992	Redds/ km 1993	Redds/ km 1994	Redds/ km 1995	Redds/ km 2000	Redds/ km 2001
Eltapom Creek	1.3	14.0	6.2	10.0	13.5	19.2	2.5	8.72	3.84
Pelletreau Creek	0.8	0	2.5	0	0	1.7	0.9	0	0
E.F. Hayfork Creek	7.4	4.3	2.7	0.6	0.4	2.2	0.3	0	1.47
Potato Creek	2.4	2.1	0	0.4	0.4	0.9	0	0.99	0.50
Tule Creek	2.7	9.5	0.5	2.2	1.6	2.4	0.4	1.9	0.48
Plummer Creek	3.4	NS	6.6	7.9	5.0	6.1	2.1	0.97	0 ₂
Rattlesnake Creek	10.6	2.6	0.8	1.8	0.4	0.8	0.1	0.74	0.31
Smoky Creek	2.4	6.6	5.0	5.9	1.9	1.9	2.1	1.63	0.73

¹ Approximate distance surveyed; in some years survey distance may have differed by 5-10%.

² Poor conditions and less numerous surveys portray fewer redds or fish than most likely were present.

Problems are commonplace and often complicate and sometimes prevent redd surveys from occurring. Some of these problems create suspicion or bias within the data, while some prevent the proper coverage of a selected tributary.

Two creeks, Big French and EFNF Trinity were dropped this year due to seasonal flow and size problems. Five adult steelhead were observed in EFNF in early March, when crews had to snorkel the river at high flows. During a later safety meeting, we decided the quality of data gathered on EFNF was not worth the high safety risks involved. In both of these creeks, surveys cannot be safely conducted when the majority of steelhead are migrating upstream and spawning in the watershed. Two other creeks, Smoky and Plummer, were not surveyed the requisite two or more times due to weather and flow

conditions. Overall, these surveys indicate that steelhead spawners were less numerous in Trinity River tributaries than in previous years.

One primary problem that affects sample design, as well as proper and even coverage, is access. Most of the Trinity basin is composed of rugged mountainous terrain with little road coverage. Roads that do exist are often poorly maintained logging roads, which rarely lead to the confluence of a selected tributary. Some tributaries lie within wilderness areas, where no roads exist, and hiking in to survey is the only possibility. Access problems are further compounded by extreme winter conditions such as snow storms. Some tributaries, such as the East Fork of the South Fork are not accessible by road until late April due to heavy snowpack.

Permission for access to private property is another possible complication affecting spawning surveys. Even though the riverbeds of all navigable rivers are held in the public trust by the State of California, it is the Department's policy to ascertain permission to cross private property, even when questioned tributaries are clearly navigable. Most of the Trinity basin is composed of US Forest Service and Bureau of Land Management property, but small portions do exist which are privately owned, either by private landowners or large timber companies. Permission had to be ascertained for access to all streams that flow through private property. Only one tributary, Rush Creek, had to be dropped because of lack of/denial of permission. Redding Creek also had refusals to be surveyed, but this was overcome by getting permission from the adjacent landowner. Up to recently there was one remaining landowner refusing access through his property with no adjacent landowner to ascertain permission from. This resulted in the crew having to turn around at his property line and backtrack approximately 400 meters and then leapfrog his 400 meters of property, creating a small hole in coverage. This problem was alleviated mid-season when we discovered the landowner had died. The executor of the will granted Fish and Game temporary permission to access and offered the property for sale to the State of California. Sierra Pacific Industries was extremely helpful; they granted permission to conduct surveys on five different tributaries crossing their property.

Another problem facing the spawning survey is selection of the area to be surveyed within a selected tributary. In a perfect world, every tributary would be surveyed from confluence to a complete upstream barrier to migration. In reality, many barriers are not well documented; or were documented when popular knowledge identified log jams as impassable. All but four tributaries were surveyed to a complete barrier. E.F.N.F. of the Trinity River was surveyed to an unnamed tributary upstream of the East Branch. Big French Creek was surveyed to Willow Gulch, Redding Creek was Byron Gulch, and E.F. Hayfork Creek was surveyed to a different Byron Gulch. These cut-off points were selected based on extreme distance from access or lack of habitat.

Historically, only one or two passes was made of each tributary to quantify spawning. This could have possibly have left some redds unidentified, especially if surveys were late in the season, or interrupted by high flow events. During the 2001 survey, a minimum of two, and sometimes three passes of each tributary were made. Surveys are

all offset by at least one month, in order to provide the best temporal coverage of the season as possible.

Recommendations

I would recommend the following additions and changes to the redd survey to gain a more complete understanding of steelhead spawning in the Trinity River basin.

A current survey of barriers to anadromous fish passage should be completed for the entire Trinity basin. Most habitat data available are incomplete and scattered around multiple agencies. Many tributaries that flow through private property have never been surveyed, so all useable habitat estimates are made from aerial photographs and topographic maps. Also, much of the data is antiquated; the definition of what constitutes a barrier to steelhead migration has dramatically changed within the last 10-20 years. In the 1970's, it was popular knowledge among fisheries professionals that a six foot log jam was a barrier to anadromous salmonids. We now know that this is simply untrue, it is possible for a perfectly conditioned adult steelhead to jump in excess of fifteen vertical feet (WDFW, 1999).

Surveys should be conducted more frequently to identify redds that are possibly covered during spring high flow events. Michael Dean, of CDFG, suggested conducting surveys weekly for maximum coverage (pers comm, 1999). During year 2000 surveys, South Fork of Indian Creek and Dutch Creek were chosen to be surveyed at double the sampling interval, because of its close proximity to the Weaverville office, to examine if more frequent surveys would identify more redds. This additional sampling effort was terminated when neither creek produced redds by mid-season.

There should be standardization of survey distances of the same creek between years. Whenever possible, tributaries were surveyed from their confluence to an upstream migrational barrier. Year to year variation in surveyed distances creates problems when making comparisons between years. When distances are not equal only redd density is comparable, and problems could arise with quality of included habitat between years.

I have several logistical recommendations that I believe will make surveys easier and more consistent between years and crew members. Crews should camp whenever possible to reduce commute time on multiple day surveys. Quad-runner all-terrain vehicles could be used to access certain tributaries that are otherwise inaccessible due to snow. Yearly training should be conducted for the crew for Swiftwater Rescue, Garmin global positioning system use, and steelhead redd identification. Finally, I would recommend daily data entry and downloading of map coordinates to simplify data storage and processing.

Literature Cited

- CDFG. 1991-1996. Annual Reports Trinity Basin Salmon and Steelhead Monitoring Project 1989-1995 seasons. Inland Fisheries Division.
- CDFG. 1998. Strategic plan for management of Klamath Mountains Province steelhead trout. Prepared for the National Marine Fisheries Service by the Resources Agency, 15 pp.
- Elms-Cockrum, T.J. 1997. Salmon spawning ground surveys, 1996. Idaho Department of Fish and Game Report 97-25, Boise, Idaho.
- LaFaunce, D.A. 1964. A Steelhead Spawning Survey of the Upper Trinity River System, 1964. Marine Resources Administrative Report No. 65-4. California Department of Fish and Game, Region 1, Inland Fisheries Branch.
- Magnusson, W.E., G.J. Caughley, and G.C. Grigg. 1978. A double-survey estimate of population size from incomplete. *Journal of Wildlife Management* 42:174-176.
- Riemen, B.E., and D.L. Myers. 1997. Use of redd counts to detect trends in bull trout (*Salvelinus confluentus*) populations. *Conservation Biology* 11:1015-1018.
- Rogers, D.W. 1971. A Steelhead Spawning Survey of The Tributaries of the Upper Trinity River and Upper Hayfork Creek Drainage. California Department of Fish and Game, Region 1, Inland Fisheries Branch.
- WDFW. 1999. Fish Passage Design at Road Culverts: a design manual for fish crossing at road culverts. Environmental Engineering Services.